



ACC.14

TCT@ACC-12 | innovation in intervention

A1374

JACC April 1, 2014

Volume 63, Issue 12



## Prevention

**=EFFICACY AND SAFETY OF BOCOCIZUMAB (RN316/PF-04950615), A MONOCLONAL ANTIBODY AGAINST PROPROTEIN CONVERTASE SUBTILISIN/KEXIN TYPE 9 IN STATIN-TREATED HYPERCHOLESTEROLEMIC SUBJECTS: RESULTS FROM A RANDOMIZED, PLACEBO-CONTROLLED, DOSE-RANGING STUDY (NCT: 01592240)**

Poster Contributions

Hall C

Sunday, March 30, 2014, 9:45 a.m.-10:30 a.m.

Session Title: Prevention: Familial Hypercholesterolemia, Novel Therapies and Cardiovascular Risk

Abstract Category: 20. Prevention: Clinical

Presentation Number: 1183-129

Authors: *Christie M. Ballantyne, Joel Neutel, Anne Cropp, William Duggan, Ellen Wang, David Plowchalk, Kevin Sweeney, Nitin Kaila, John Vincent, Harold Bays, Baylor College of Medicine and Houston Methodist DeBakey Heart and Vascular Center, Houston, TX, USA, Pfizer Inc., Groton and New York, NY, USA*

**Background:** Bococizumab (BOCO) is a humanized monoclonal antibody that binds to proprotein convertase subtilisin/kexin type 9 and is a therapeutic approach for reducing LDL-C levels in hypercholesterolemic (HC) patients. The objective of this study was to evaluate various BOCO subcutaneous (SC) doses administered every 2 weeks (Q14d) or monthly (Q28d).

**Methods:** This was a 24-week, multicenter, randomized, double-blind, placebo-controlled trial in statin-treated subjects aged  $\geq 18$  yrs with a diagnosis of HC and LDL-C  $\geq 80$  mg/dL. Subjects were randomized to Q14d SC placebo, BOCO 50 mg, 100 mg, or 150 mg; or Q28d placebo, BOCO 200 mg or 300 mg. Doses were reduced if LDL-C readings were  $\leq 25$  mg/dL. The primary analysis was the placebo-adjusted treatment difference for the change from baseline in LDL-C at Week 12.

**Results:** BOCO significantly reduced LDL-C across all doses (Table). Up to 44% of subjects in the BOCO groups had their dose reduced. Model-based simulation that assumed no dose reductions predicted greater LDL-C lowering than the primary endpoint, which included subjects having doses reduced. AEs were similar across placebo and BOCO groups. Few subjects discontinued treatment due to treatment-related AEs (Table).

**Conclusions:** BOCO 150 mg Q14d significantly reduced LDL-C by 53 mg/dL vs. placebo at Week 12, inclusive of the protocol-directed dose reductions in 35% of subjects. Modeling predicted greater LDL-C reduction in absence of BOCO dose reduction. The Q14d regimen is being evaluated in larger trials.

	Placebo Q14d (n=49)	BOCO 50 mg Q14d (n=50)	BOCO 100 mg Q14d (n=51)	BOCO 150 mg Q14d (n=49)	Placebo Q28d (n=51)	BOCO 200 mg Q28d (n=50)	BOCO 300 mg Q28d (n=51)
Baseline LDL-C, mg/dL	108.7 $\pm$ 31.5	107.9 $\pm$ 20.2	113.4 $\pm$ 25.7	105.8 $\pm$ 18.0	118.8 $\pm$ 44.8	105.7 $\pm$ 23.2	104.7 $\pm$ 22.1
Subjects with dose reduction	0	0	16%	35%	0	44%	39%
Absolute change from baseline in LDL-C at Week 12, mg/dL	-2.8 $\pm$ 29.2	-35.4 $\pm$ 26.6	-52.3 $\pm$ 31.3	-54.2 $\pm$ 27.0	-1.3 $\pm$ 37.2	-21.3 $\pm$ 28.0	-38.3 $\pm$ 41.3
Adjusted change from baseline in LDL-C at Week 12, versus placebo,* mg/dL	---	-34.3 (-45.1, -23.5)	-45.1 (-55.9, -34.2)	-53.4 (-64.1, -42.7)	---	-27.6 (-40.5, -14.7)	-44.9 (-57.7, -32.1)
Model predicted placebo-adjusted LDL-C change from baseline at Week 12 assuming no dose reduction,** mg/dL	---	-36.3 (-42.0, -30.3)	-60.6 (-67.1, -55.5)	-72.1 (-77.5, -67.0)	---	-40.3 (-47.2, -34.3)	-55.7 (-61.6, -48.0)
Subjects with treatment-related SAEs	0	0	0	2%	0	0	0
Discontinuation of treatment due to treatment-related AEs	0	2%	0	8%	0	0	4%

Values are mean  $\pm$  SD, %, or n=number treated, also see \* and \*\* below. BOCO, bococizumab; LDL-C, low-density lipoprotein cholesterol; AE, adverse event; SAE, serious AE.

\*Mean (95% confidence interval), calculated using a repeated measures model for longitudinal data.

\*\*Population PK/PD model predictions of median absolute LDL-C change from baseline (2.5th and 97.5th percentile) for 100 trials (50 subjects/arm) using baseline demographics observed in this study.